



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Number of Terms.	Mid-Term or Pair of Mid-Terms.	The Series.
1	15	15
3	5	4+5+6
5	3	1+2+3+4+5
15	1	-6-5-4-3-2-1+0+1+2+3+4+5+6+7+8
2	7,8	7+8
6	2,3	0+1+2+3+4+5
10	1,2	-3-2-1+0+1+2+3+4+5+6
30	0,1	-14-13- . . . +0+1 . . . +14+15

NOTE ON PRIME NUMBERS.

By DERRICK N. LEHMER, University of California.

It is a well known theorem that it is possible to find an arbitrarily great number of consecutive composite numbers. This appears from the values which the expression $n!+r$ takes for $r=2, 3, \dots, n$. This theorem furnishes an interesting proof of the theorem that the number of primes less than or equal to x is not determined by a function of x which is a polynomial in x of finite degree. For if $f(x)$ were such a function of degree n , then for $x=(n+2)!+r$, $f(x)$ must keep the same value for $r=2, 3, 4, \dots, n+2$. If this value is k , then $f(x)-k=0$ is an equation of degree n with $n+1$ roots, which is impossible.

DEPARTMENTS.

SOLUTIONS OF PROBLEMS.

ALGEBRA.

362. Proposed by JAMES F. LAWRENCE, Stillwater, Okla.

Show that the number of solutions in positive integers, zero included, of the equation $x+2y+3z=6n$, is $3n^2+3n+1$.

Solution by W. J. GREENSTREET, M. A., Editor of the Mathematical Gazette, Burghfield, England.

$x+2y+3z=6n$. z may have any value from 0 to $2n$, inclusive.

Hence we may assign to it any even or odd value from 0 to $2n$, inclusive.